

l99\_jordan  
(TMNxpQnkGP1B9tPTnrQcC24vA3xVCffvwy)

October 27, 2020

Let  $k17\_euclid : \iota \Rightarrow \iota$  be given. Let  $k19\_euclid : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k1\_real\_1 : \iota \Rightarrow \iota$  be given. Let  $np\_3 : \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k18\_euclid : \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_xcmplx\_0 : \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow ((k17\_euclid (k19\_euclid X0 X1) = X0) \wedge (k18\_euclid (k19\_euclid X0 X1) = X1))) \quad (1)$$

Assume the following.

$$((v2\_xxreal\_0 np\_3) \wedge (m2\_subset\_1 np\_3 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_3 k5\_numbers) \wedge (m1\_subset\_1 np\_3 k1\_numbers)) \quad (2)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (k1\_real\_1 X0 = k4\_xcmplx\_0 X0) \quad (3)$$

Assume the following.

$$m1\_subset\_1 k6\_numbers k1\_numbers \quad (4)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow ((v1\_xcmplx\_0 (k4\_xcmplx\_0 X0)) \wedge (v1\_xreal\_0 (k4\_xcmplx\_0 X0))) \quad (5)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (6)$$

**Theorem 1**  $k17\_euclid (k19\_euclid k6\_numbers (k1\_real\_1 np\_3)) = k6\_numbers.$