

t100_finseq_2 (TMHGh- hYtXr66DUZztFUfYuxFjHuYouFTWYw)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k3_finseq_2 : \iota \Rightarrow \iota$ be given. Let $k13_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (k4_finseq_2 np_2 X0 = ReplSep2 \\ (toset (\lambda X1 : \iota. m1_subset_1 X1 X0)) (\lambda X1 : \iota. toset (\lambda X2 : \\ \iota. m1_subset_1 X2 X0)) (\lambda X1 : \iota. \lambda X2 : \iota. True) (\lambda X1 : \\ \iota. \lambda X2 : \iota. k10_finseq_1 X1 X2)) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (2)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_2 X1 X0) \Rightarrow (\forall X2. (m2_finseq_2 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Leftrightarrow(m1_finseq_1 X1 X0) \quad (5)$$

Assume the following.

$$\forall X0.k9_finseq_1 X0 = k5_finseq_1 X0 \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.k3_finseq_2 X0 = k13_finseq_1 X0 \quad (8)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow (k3_finseq_1 X0 = k1_card_1 X0) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (11)$$

Assume the following.

$$\forall X0.m1_finseq_2 (k3_finseq_2 X0) X0 \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.k10_finseq_1 X0 X1 = k7_finseq_1 (k9_finseq_1 X0) (k9_finseq_1 X1) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(v3_card_1 X1 X0)\Leftrightarrow(k1_card_1 X1 = X0) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.k4_tarski X0 X1 = k2_tarski (k2_tarski X0 X1) (k1_tarski X0) \quad (15)$$

Assume the following.

$$\forall X0. k5_finseq_1 X0 = k1_tarSKI (k4_tarSKI np_1 X0) \quad (16)$$

Assume the following.

$$\begin{aligned} \forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. k4_finseq_2 X0 X1 = ReplSep \\ (toset (\lambda X2 : \iota. m2_finseq_2 X2 X1 (k3_finseq_2 X1))) (\lambda X2 : \\ \iota. k3_finseq_1 X2 = X0) (\lambda X2 : \iota. X2))) \end{aligned} \quad (17)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k13_finseq_1 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (m2_finseq_1 X2 X0)) \quad (18)$$

Assume the following.

$$\forall X0. \forall X1. k2_tarSKI X0 X1 = k2_tarSKI X1 X0 \quad (19)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (20)$$

Theorem 1

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v3_card_1 X1 np_2) \wedge \\ (m2_finseq_1 X1 X0)) \Rightarrow (\exists X2. (m1_subset_1 X2 X0) \wedge (\exists X3. \\ (m1_subset_1 X3 X0) \wedge (X1 = k10_finseq_1 X2 X3)))) \end{aligned}$$