

t100_seq_4

(TMbwh7VYjVU7uqA8XKsGR4K4e7GAH2Rk7Wp)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_numbers : \iota$ be given. Let $k14_seq_4 : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k22_seq_4 : \iota \Rightarrow \iota$ be given. Let $k15_seq_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k19_seq_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & (m2_finseq_2 X1 k2_numbers (k14_seq_4 X0)) \Rightarrow (\forall X2.(m2_finseq_2 \\ & X2 k2_numbers (k14_seq_4 X0)) \Rightarrow (r1_xxreal_0 (k22_seq_4 (k19_seq_4 \\ & X0 X1 X2)) (k9_binop_2 (k22_seq_4 X1) (k22_seq_4 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & (m2_finseq_2 X1 k2_numbers (k14_seq_4 X0)) \Rightarrow (\forall X2.(m2_finseq_2 \\ & X2 k2_numbers (k14_seq_4 X0)) \Rightarrow (X1 = k19_seq_4 X0 (k15_seq_4 X0 X1 \\ & X2) X2))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 (k2_xcmplx_0 X1 X2)) \Leftrightarrow (r1_xxreal_0 \\ & (k6_xcmplx_0 X0 X1) X2)))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Leftrightarrow(m1_subset_1 X2 X1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0)\Rightarrow(\forall X2.(m2_finseq_2 X2 X0 X1)\Leftrightarrow(m1_subset_1 X2 X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Leftrightarrow(m1_finseq_1 X1 X0) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(k9_binop_2 X0 X1 = k2_xcmplx_0 X0 X1) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_subset_1 X0 k5_numbers)\wedge((m1_subset_1 X1 (k14_seq_4 X0))\wedge(m1_subset_1 X2 (k14_seq_4 X0))))\Rightarrow(k15_seq_4 X0 X1 X2 = k1_valued_1 X1 X2) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(k10_binop_2 X0 X1 = k6_xcmplx_0 X0 X1) \quad (9)$$

Assume the following.

$$\exists X0.(v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge((\neg v1_xboole_0 X0)\wedge(v1_finset_1 X0)))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0)\Rightarrow(\forall X2.(m2_finseq_2 X2 X0 X1)\Rightarrow(m2_finseq_1 X2 X0)) \quad (11)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (12)$$

Assume the following.

$$\forall X0.(m1_finseq_1 X0 k2_numbers)\Rightarrow(m1_subset_1 (k22_seq_4 X0) k1_numbers) \quad (13)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((m1_subset_1 X0 k5_numbers)\wedge \\ & ((m1_subset_1 X1 (k14_seq_4 X0))\wedge(m1_subset_1 X2 (k14_seq_4 X0))))\Rightarrow \\ & (m2_finseq_2 (k15_seq_4 X0 X1 X2) k2_numbers (k14_seq_4 X0)) \end{aligned} \quad (14)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k5_numbers)\Rightarrow(m1_finseq_2 (k14_seq_4 X0) k2_numbers) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(k9_binop_2 X0 X1 = k9_binop_2 X1 X0) \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((m1_subset_1 X0 k5_numbers)\wedge \\ & ((m1_subset_1 X1 (k14_seq_4 X0))\wedge(m1_subset_1 X2 (k14_seq_4 X0))))\Rightarrow \\ & (k15_seq_4 X0 X1 X2 = k15_seq_4 X0 X2 X1) \end{aligned} \quad (17)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow((v1_xboole_0 X1)\wedge((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0)))) \quad (18)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (19)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_xboole_0 X1)) \quad (20)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers)\Rightarrow(\forall X1. \\ & (m2_finseq_2 X1 k2_numbers (k14_seq_4 X0))\Rightarrow(\forall X2.(m2_finseq_2 \\ & X2 k2_numbers (k14_seq_4 X0))\Rightarrow(r1_xxreal_0 (k10_binop_2 (k22_seq_4 \\ & X1) (k22_seq_4 X2)) (k22_seq_4 (k15_seq_4 X0 X1 X2)))))) \end{aligned}$$