

t101_flang_2
(TMdbV3J7AJgbAgNSJ26pj6LLYPSBMnrwe8r)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_flang_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_flang_1 : \iota \Rightarrow \iota$ be given. Let $k3_catalan2 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k8_afinsq_1 X1))) \Rightarrow ((X0 \in k2_flang_2 X1 X2) \Leftrightarrow ((X0 = k2_flang_1 X1) \vee \\ & (X0 \in X2))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k3_catalan2 \\ & X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k3_catalan2 \\ & X0))) \Rightarrow ((k2_flang_1 X0 \in X1) \Rightarrow ((r1_tarski X2 (k6_flang_1 X0 X2 X1)) \wedge \\ & (r1_tarski X2 (k6_flang_1 X0 X1 X2)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0. k3_catalan2 X0 = k8_afinsq_1 X0 \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & X0))) \Rightarrow (m1_subset_1 (k2_flang_2 X0 X1) (k1_zfmisc_1 (k8_afinsq_1 \\ & X0))) \end{aligned} \tag{4}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k8_afinsq_1 \\ & X0))) \Rightarrow ((r1_tarski X1 (k6_flang_1 X0 X1 (k2_flang_2 X0 X2))) \wedge (r1_tarski \\ & X1 (k6_flang_1 X0 (k2_flang_2 X0 X2) X1)))) \end{aligned}$$