

t101_funct_1
(TMXR1yCxQ96oHmHyA4V9aXfM2siM82FDrF)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. \\ & ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 (k3_relat_1 \\ & X2 X1)) \Leftrightarrow ((X0 \in k9_xtuple_0 X2) \wedge (k1_funct_1 X2 X0 \in k9_xtuple_0 X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & (X2 = k7_relat_1 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow (\exists X4. (X4 \in k9_xtuple_0 \\ & X0) \wedge ((X4 \in X1) \wedge (X3 = k1_funct_1 X0 X4)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((\\ & v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. (r1_tarski X2 (k9_xtuple_0 \\ & (k3_relat_1 X0 X1))) \Leftrightarrow ((r1_tarski X2 (k9_xtuple_0 X0)) \wedge (r1_tarski \\ & (k7_relat_1 X0 X2) (k9_xtuple_0 X1)))))) \end{aligned}$$