

t101_funct_4
(TMXRWdyi9k9ZmGqDfhiL7dSx3bja86bvELC)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k6_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. X0 \in k9_xtuple_0 (k16_funcop_1 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k1_funct_1 (k16_funcop_1 X0 X1) X0 = X1 \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (r1_tarski (k10_xtuple_0 X0) (k10_xtuple_0 (k1_funct_4 X1 X0)))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 (k3_relat_1 X2 X1)) \Rightarrow (k1_funct_1 (k3_relat_1 X2 X1) X0 = k1_funct_1 X1 (k1_funct_1 X2 X0)))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 (k3_relat_1 X2 X1)) \Leftrightarrow ((X0 \in k9_xtuple_0 X2) \wedge (k1_funct_1 X2 X0 \in k9_xtuple_0 X1)))) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \wedge ((v1_relat_1 X1) \wedge (v1_funct_1 X1))) \Rightarrow ((v1_relat_1 (k3_relat_1 X0 X1)) \wedge (v1_funct_1 (k3_relat_1 X0 X1))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 (k16_funcop_1 X0 X1))\wedge(v1_funct_1 (k16_funcop_1 X0 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.v1_relat_1 (k3_relat_1 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.\forall X2. k6_funct_4 X0 X1 X2 = k1_funct_4 X0 (k3_relat_1 X0 (k16_funcop_1 X1 X2))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow (X2 \in X1)) \quad (10)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.(X1 = k10_xtuple_0 X0)\Leftrightarrow(\forall X2.(X2 \in X1)\Leftrightarrow(\exists X3.(X3 \in k9_xtuple_0 X0)\wedge(X2 = k1_funct_1 X0 X3)))) \quad (11)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.\forall X2. (X1 \in k10_xtuple_0 X0)\Rightarrow(X2 \in k10_xtuple_0 (k6_funct_4 X0 X1 X2)))$$