

t102_mesfunc6 (TM- FWAAVZ7hHqAMHk1iNwtaRv3c7RcLBG6AD)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $v10_valued_0 : \iota \Rightarrow o$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $v4_measure1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r3_mesfunc6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k26_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_mesfunc6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_extreal1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_measure6 : \iota \Rightarrow \iota$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_mesfunc5 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_mesfunc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_mesfunc5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_mesfunc5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge \\ & m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ & \forall X2. (m1_subset_1 X2 k1_numbers) \Rightarrow (r2_relset_1 X0 k7_numbers \\ & (k1_mesfunc5 X0 (k26_valued_1 X0 k1_numbers X1 X2)) (k6_mesfunc1 \\ & X0 (k1_mesfunc5 X0 X1) X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\ & X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))))) \Rightarrow \\ & (\forall X3. ((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k7_numbers)))) \Rightarrow (\forall X4. (m1_subset_1 X4 k1_numbers) \Rightarrow (\\ & (r1_mesfunc5 X0 X1 X2 X3) \Rightarrow ((r1_mesfunc5 X0 X1 X2 (k6_mesfunc1 X0 \\ & X3 X4)) \wedge (k7_mesfunc5 X0 X1 X2 (k6_mesfunc1 X0 X3 X4) = k1_extreal1 \\ & (k1_measure6 X4) (k7_mesfunc5 X0 X1 X2 X3)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((r2_relset_1 X0 X1 X2 X3)\Leftrightarrow(X2 = X3)) \quad (3)$$

Assume the following.

$$v3_membered k1_numbers \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(((v1_relat_1 X1)\wedge((v4_relat_1 X1 X0)\wedge((v5_relat_1 X1 k7_numbers)\wedge(v1_funct_1 X1))))\wedge(m1_subset_1 X2 k1_numbers)))\Rightarrow((v1_funct_1 (k6_mesfunc1 X0 X1 X2))\wedge(m1_subset_1 (k6_mesfunc1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v3_membered X1)\wedge(((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\wedge(v1_xreal_0 X3)))\Rightarrow((v1_funct_1 (k26_valued_1 X0 X1 X2 X3))\wedge(m1_subset_1 (k26_valued_1 X0 X1 X2 X3) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow((v1_funct_1 (k1_mesfunc5 X0 X1))\wedge(m1_subset_1 (k1_mesfunc5 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \quad (7)$$

Assume the following.

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow(k1_mesfunc5 X0 X1 = X1)) \quad (8)$$

Assume the following.

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((\neg v1_xboole_0 X1)\wedge((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge((v1_funct_2 X2 X1 k7_numbers)\wedge((v10_valued_0 X2)\wedge((v6_supinf_2 X2)\wedge((v4_measure1 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers))))))))\Rightarrow(\forall X3.((v1_funct_1 X3)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow((r3_mesfunc6 X0 X1 X2 X3)\Leftrightarrow(r1_mesfunc5 X0 X1 X2 (k1_mesfunc5 X0 X3)))))) \quad (9)$$

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\
& X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
& X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
& (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
& X0 k1_numbers)))) \Rightarrow (k1_mesfunc6 X0 X1 X2 X3 = k7_mesfunc5 X0 X1 X2 \\
& (k1_mesfunc5 X0 X3))))))
\end{aligned} \tag{10}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \tag{11}$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \tag{12}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \tag{13}$$

Theorem 1

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\
& X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
& X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
& (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
& X0 k1_numbers)))) \Rightarrow (\forall X4. (m1_subset_1 X4 k1_numbers) \Rightarrow (\\
& (r3_mesfunc6 X0 X1 X2 X3) \Rightarrow ((r3_mesfunc6 X0 X1 X2 (k26_valued_1 X0 \\
& k1_numbers X3 X4) \wedge (k1_mesfunc6 X0 X1 X2 (k26_valued_1 X0 k1_numbers \\
& X3 X4) = k1_extreal1 (k1_measure6 X4) (k1_mesfunc6 X0 X1 X2 X3))))))
\end{aligned}$$