

t102_scmfsa_2 (TMQsjyC-
PagQ8QTVrXPDj6fzmJF7Q9qKAV1L)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_ami_2 : \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_scmfsa_2 : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_memstr_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $k4_struct_0 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_memstr_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_extpro_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_memstr_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $l1_extpro_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $l1_memstr_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $l1_compos_1 : \iota \Rightarrow o$ be given. Let $v1_setfam_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v5_funct_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_memstr_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((v1_ami_2 X0) \wedge (m1_subset_1 X0 (u1_struct_0 k1_scmfsa_2))) \Rightarrow (X0 \neq k4_struct_0 k1_scmfsa_2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \quad (3)$$

Assume the following.

$$\neg v1_xboole_0 np_3 \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(\quad (7)$$

$$k1_relset_1 X0 X1 = k9_xtuple_0 X1)$$

Assume the following.

$$(v3_memstr_0 k1_scmf_sa_2 np_3)\wedge(v1_extpro_1 k1_scmf_sa_2 np_3) \quad (8)$$

Assume the following.

$$(\neg v2_struct_0 k1_scmf_sa_2)\wedge((v2_memstr_0 k1_scmf_sa_2 np_3)\wedge \quad (9)$$

$$(v1_extpro_1 k1_scmf_sa_2 np_3))$$

Assume the following.

$$\forall X0.\forall X1.(l1_extpro_1 X1 X0)\Rightarrow((l1_memstr_0 X1 X0)\wedge \quad (10)$$

$$(l1_compos_1 X1))$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_setfam_1 X0)\wedge(((\neg v2_struct_0 \quad (11)$$

$$X1)\wedge((v2_memstr_0 X1 X0)\wedge((v3_memstr_0 X1 X0)\wedge(l1_memstr_0 X1$$

$$X0))))\wedge(v7_ordinal1 X2))\Rightarrow((v1_relat_1 (k7_memstr_0 X0 X1 X2))\wedge$$

$$((v4_relat_1 (k7_memstr_0 X0 X1 X2) (u1_struct_0 X1))\wedge((v1_funct_1$$

$$(k7_memstr_0 X0 X1 X2))\wedge(v5_funct_1 (k7_memstr_0 X0 X1 X2) (k2_memstr_0$$

$$X0 X1))))))$$

Assume the following.

$$(v1_extpro_1 k1_scmf_sa_2 np_3)\wedge(l1_extpro_1 k1_scmf_sa_2 np_3) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.k16_funcop_1 X0 X1 = k7_funcop_1 (k1_tarski \quad (13)$$

$$X0) X1$$

Assume the following.

$$\forall X0.\forall X1.k2_funcop_1 X0 X1 = k2_zfmisc_1 X0 (k1_tarski \quad (14)$$

$$X1)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarSKI X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (15)$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_setfam_1 X0) \Rightarrow (\forall X1.((\neg v2_struct_0 X1) \wedge \\ ((v2_memstr_0 X1 X0) \wedge (v3_memstr_0 X1 X0) \wedge (l1_memstr_0 X1 X0)))) \Rightarrow \\ (\forall X2.(v7_ordinal1 X2) \Rightarrow (k7_memstr_0 X0 X1 X2 = k16_funcop_1 \\ (k4_struct_0 X1) X2))) \end{aligned} \quad (16)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (17)$$

Assume the following.

$$\forall X0.((\neg v1_xboole_0 X0) \wedge (v7_ordinal1 X0)) \Rightarrow ((\neg v1_xboole_0 X0) \wedge ((v7_ordinal1 X0) \wedge (\neg v1_setfam_1 X0))) \quad (18)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.((v1_ami_2 \\ X1) \wedge (m1_subset_1 X1 (u1_struct_0 k1_scmfSA_2))) \Rightarrow (\neg X1 \in k1_relset_1 \\ (u1_struct_0 k1_scmfSA_2) (k7_memstr_0 np_3 k1_scmfSA_2 X0))) \end{aligned}$$