

t102\_xboole\_1  
(TMbpdJ7gbTq7ESK4Hh5VoxjTsS5Jx3DX7Fg)

October 27, 2020

Let  $k4\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. k4\_xboole\_0 X0 (k4\_xboole\_0 X1 X2) = k2\_xboole\_0 (k4\_xboole\_0 X0 X1) (k3\_xboole\_0 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. k3\_xboole\_0 (k3\_xboole\_0 X0 X1) X2 = k3\_xboole\_0 X0 (k3\_xboole\_0 X1 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k5\_xboole\_0 X0 X1 = k4\_xboole\_0 (k2\_xboole\_0 X0 X1) (k3\_xboole\_0 X0 X1) \quad (3)$$

**Theorem 1**

$$\forall X0. \forall X1. \forall X2. k4\_xboole\_0 X0 (k5\_xboole\_0 X1 X2) = k2\_xboole\_0 (k4\_xboole\_0 X0 (k2\_xboole\_0 X1 X2)) (k3\_xboole\_0 (k3\_xboole\_0 X0 X1) X2)$$