

t103\_flang\_2

(TMLMg9aHWqVxhqCQPkqvCDSvTV23Qyshu51)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k8\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k2\_flang\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_flang\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_catalan2 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow ((k2\_flang\_2 X0 X1 = X1) \Leftrightarrow (k2\_flang\_1 X0 \in X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow (\forall X2. (v7\_ordinal1 X2) \Rightarrow ((k2\_flang\_1 X0 \in X1) \Rightarrow ((r1\_xxreal\_0 X2 k6\_numbers) \vee (r1\_tarski X1 (k7\_flang\_1 X0 X1 X2)))))) \quad (2)$$

Assume the following.

$$\forall X0. k3\_catalan2 X0 = k8\_afinsq\_1 X0 \quad (3)$$

**Theorem 1**

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow (\forall X2. (v7\_ordinal1 X2) \Rightarrow ((k2\_flang\_1 X0 \in X1) \Rightarrow ((r1\_xxreal\_0 X2 k6\_numbers) \vee (r1\_tarski (k2\_flang\_2 X0 X1) (k7\_flang\_1 X0 X1 X2))))))$$