

t104\_chord (TMPoRz-  
toCi6J2MPJDn9KxmhwaG8DRB2Lq2Q)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $v1\_glib\_000 : \iota \Rightarrow o$  be given. Let  $v2\_glib\_000 : \iota \Rightarrow o$  be given. Let  $m3\_chord : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k5\_card\_1 : \iota \Rightarrow \iota$  be given. Let  $k6\_glib\_000 : \iota \Rightarrow \iota$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $v2\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & ((v2\_funct\_1 X0) \Leftrightarrow (k5\_card\_1 (k10\_xtuple\_0 X0) = k3\_finseq\_1 X0)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 \\ & X0) \wedge ((v1\_finset\_1 X0) \wedge ((v1\_glib\_000 X0) \wedge (v2\_glib\_000 X0)))))) \Rightarrow \\ & (\forall X1.(m3\_chord X1 X0) \Rightarrow (m2\_finseq\_1 X1 (k6\_glib\_000 X0))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Rightarrow ((v1\_funct\_1 X1) \wedge ( \\ & (v1\_finseq\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers \\ & X0)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 \\ & X0) \wedge ((v1\_finset\_1 X0) \wedge ((v1\_glib\_000 X0) \wedge (v2\_glib\_000 X0)))))) \Rightarrow \\ & (\forall X1.(m2\_finseq\_1 X1 (k6\_glib\_000 X0) \Rightarrow ((m3\_chord X1 X0) \Leftrightarrow \\ & ((v2\_funct\_1 X1) \wedge (k10\_xtuple\_0 X1 = k6\_glib\_000 X0)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \end{aligned} \quad (5)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 \ k5\_numbers) \wedge ((v1\_funct\_1 \\ & X0) \wedge ((v1\_finset\_1 X0) \wedge ((v1\_glib\_000 X0) \wedge (v2\_glib\_000 X0))))) \Rightarrow \\ & (\forall X1.(m3\_chord X1 X0) \Rightarrow (k3\_finseq\_1 X1 = k5\_card\_1 (k6\_glib\_000 \\ & X0))) \end{aligned}$$