

t104_flang_2
(TMGNUAS6aRNvxGUi3hU5AjKp8vUDHckKF41)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k6_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_flang_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_flang_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0))) \Rightarrow (k2_flang_2 X0 X1 = k1_flang_2 X0 X1 k6_numbers np_1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & \quad X0))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (\forall X3. (v7_ordinal1 \\ & \quad X3) \Rightarrow (\forall X4. (v7_ordinal1 X4) \Rightarrow (k6_flang_1 X0 (k1_flang_2 \\ & \quad X0 X1 X2 X3) (k7_flang_1 X0 X1 X4) = k6_flang_1 X0 (k7_flang_1 X0 X1 \\ & \quad X4) (k1_flang_2 X0 X1 X2 X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$m1_subset_1 k1_xboole_0 k4_ordinal1 \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & X0))) \Rightarrow (\forall X2.(v7_ordinal1 X2) \Rightarrow (k6_flang_1 X0 (k2_flang_2 \\ & X0 X1) (k7_flang_1 X0 X1 X2) = k6_flang_1 X0 (k7_flang_1 X0 X1 X2) (\\ & k2_flang_2 X0 X1))) \end{aligned}$$