

t104_seq_4
(TMSJ3iSx7cTaAYQsfCe5QdSxD3MFyQcoowJ)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_numbers : \iota$ be given. Let $k14_seq_4 : \iota \Rightarrow \iota$ be given. Let $k22_seq_4 : \iota \Rightarrow \iota$ be given. Let $k19_seq_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_seq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & (m2_finseq_2 X1 k2_numbers (k14_seq_4 X0)) \Rightarrow (k22_seq_4 (k18_seq_4 \\ & X0 X1) = k22_seq_4 X1)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & (m2_finseq_2 X1 k2_numbers (k14_seq_4 X0)) \Rightarrow (\forall X2.(m2_finseq_2 \\ & X2 k2_numbers (k14_seq_4 X0)) \Rightarrow (k18_seq_4 X0 (k19_seq_4 X0 X1 X2) = \\ & k19_seq_4 X0 X2 X1))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0) \Rightarrow (\forall X2.(m2_finseq_2 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \tag{4}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{5}$$

Assume the following.

$$(\neg v1_xboole_0 \ k4_ordinal1) \wedge (v3_ordinal1 \ k4_ordinal1) \quad (6)$$

Assume the following.

$$\neg v1_xboole_0 \ k1_numbers \quad (7)$$

Assume the following.

$$m1_subset_1 \ k5_numbers \ (k1_zfmisc_1 \ k1_numbers) \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((m1_subset_1 \ X0 \ k5_numbers) \wedge \\ & ((m1_subset_1 \ X1 \ (k14_seq_4 \ X0)) \wedge (m1_subset_1 \ X2 \ (k14_seq_4 \ X0)))) \Rightarrow \\ & (m2_finseq_2 \ (k19_seq_4 \ X0 \ X1 \ X2) \ k2_numbers \ (k14_seq_4 \ X0)) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0. (m1_subset_1 \ X0 \ k5_numbers) \Rightarrow (m1_finseq_2 \ (k14_seq_4 \ X0) \ k2_numbers) \quad (10)$$

Theorem 1

$$\begin{aligned} & \forall X0. (m2_subset_1 \ X0 \ k1_numbers \ k5_numbers) \Rightarrow (\forall X1. \\ & (m2_finseq_2 \ X1 \ k2_numbers \ (k14_seq_4 \ X0)) \Rightarrow (\forall X2. (m2_finseq_2 \\ & X2 \ k2_numbers \ (k14_seq_4 \ X0)) \Rightarrow (k22_seq_4 \ (k19_seq_4 \ X0 \ X1 \ X2) = \\ & k22_seq_4 \ (k19_seq_4 \ X0 \ X2 \ X1)))) \end{aligned}$$