

t105_relat_1

(TMN9xSg24HQG1BMkFQXCOP6BJBHfckYrHcV)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k6_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarSKI : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X0) \Rightarrow (v1_relat_1 (k4_xboole_0 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (v1_relat_1 (k6_relat_1 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k4_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (\neg X3 \in X1))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (\forall X2. (v1_relat_1 X2) \Rightarrow ((X2 = k6_relat_1 X0 X1) \Leftrightarrow (\forall X3. \forall X4. (k4_tarSKI X3 X4 \in X2) \Leftrightarrow ((X4 \in X0) \wedge (k4_tarSKI X3 X4 \in X1))))) \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow (k6_relat_1 (k6_subset_1 X0 X1) X2 = k6_subset_1 (k6_relat_1 X0 X2) (k6_relat_1 X1 X2))$$