

t106_xcplx_1 (TMLnEgr-
GUN1uPNFeZK8H4f8WrYq2JDWekMB)

October 27, 2020

Let $v1_xcplx_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k3_xcplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_xcplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Assume the following.

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow (k3_xcplx_0 X0 (k7_xcplx_0 np_1 X1) = k7_xcplx_0 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (k7_xcplx_0 X0 X0 = np_1)) \quad (2)$$

Theorem 1

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (k3_xcplx_0 X0 (k7_xcplx_0 np_1 X0) = np_1))$$