

## t107\_funct\_4

(TMKS2ZPHhSivMgqhqXAefAD9xVwCYhwHhTR)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_funct\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k16\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ \forall X3.(X1 \neq X3) \Rightarrow (k1\_funct\_1 (k1\_funct\_4 X0 (k16\_funcop\_1 \\ X1 X2)) X3 = k1\_funct\_1 X0 X3)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarski X0 X1) \Leftrightarrow ((r1\_tarski \\ (k9\_xtuple\_0 X0) (k9\_xtuple\_0 X1)) \wedge (\forall X2.(X2 \in k9\_xtuple\_0 \\ X0) \Rightarrow (k1\_funct\_1 X0 X2 = k1\_funct\_1 X1 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarski (k9\_xtuple\_0 X0) \\ (k9\_xtuple\_0 (k1\_funct\_4 X0 X1))) \wedge (r1\_tarski (k9\_xtuple\_0 X1) \\ (k9\_xtuple\_0 (k1\_funct\_4 X0 X1)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1.(v1\_relat\_1 (k16\_funcop\_1 X0 X1)) \wedge (v1\_funct\_1 \\ (k16\_funcop\_1 X0 X1)) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1.(((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \wedge (( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1))) \Rightarrow ((v1\_relat\_1 (k1\_funct\_4 X0 \\ X1)) \wedge (v1\_funct\_1 (k1\_funct\_4 X0 X1))) \end{aligned} \quad (5)$$

### Theorem 1

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ (\neg X1 \in k9\_xtuple\_0 X0) \Rightarrow (r1\_tarski X0 (k1\_funct\_4 X0 (k16\_funcop\_1 \\ X1 X2)))) \end{aligned}$$