

t107_member_1
(TMJn9yvLLphhV9NB9z9X8eqPpCxFaE2BCLj)

October 27, 2020

Let $v2_membered : \iota \Rightarrow o$ be given. Let $r1_tarSKI : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k14_member_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarSKI X0 X1) \wedge (r1_tarSKI X0 X2)) \Rightarrow (r1_tarSKI X0 (k3_xboole_0 X1 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarSKI (k3_xboole_0 X0 X1) X0 \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. (v2_membered X0) \Rightarrow (\forall X1. (v2_membered X1) \Rightarrow (\forall X2. \\ (v2_membered X2) \Rightarrow (\forall X3. (v2_membered X3) \Rightarrow (((r1_tarSKI \\ X0 X1) \wedge (r1_tarSKI X2 X3)) \Rightarrow (r1_tarSKI (k14_member_1 X0 X2) (k14_member_1 \\ X1 X3)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarSKI X0 X0 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (v2_membered X0) \Rightarrow (v2_membered (k3_xboole_0 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (6)$$

Theorem 1

$$\begin{aligned} \forall X0. (v2_membered X0) \Rightarrow (\forall X1. (v2_membered X1) \Rightarrow (\forall X2. \\ (v2_membered X2) \Rightarrow (r1_tarSKI (k14_member_1 X0 (k3_xboole_0 X1 \\ X2)) (k3_xboole_0 (k14_member_1 X0 X1) (k14_member_1 X0 X2)))))) \end{aligned}$$