

t107_xboole_1

(TMZPDjBxsgr1BwLp3PYo95X2iB73wUMJH36)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_xboole_0 X0 X2)) \Rightarrow (r1_tarski X0 (k4_xboole_0 X1 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (r1_tarski X0 (k4_xboole_0 X1 X2)) \Rightarrow ((r1_tarski X0 X1) \wedge (r1_xboole_0 X0 X2)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k5_xboole_0 X0 X1 = k4_xboole_0 (k2_xboole_0 X0 X1) (k3_xboole_0 X0 X1) \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. (r1_tarski X0 (k5_xboole_0 X1 X2)) \Leftrightarrow ((r1_tarski X0 (k2_xboole_0 X1 X2)) \wedge (r1_xboole_0 X0 (k3_xboole_0 X1 X2)))$$