

t108_flang_2 (TMGTfjTm- mQm3nCnAJk2rTzjo7PKPykBihhv)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $k6_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_flang_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_flang_1 : \iota \Rightarrow \iota$ be given. Let $k3_catalan2 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0))) \Rightarrow ((k2_flang_2 X0 (k8_flang_1 X0 X1) = k8_flang_1 X0 X1) \wedge (k8_flang_1 X0 (k2_flang_2 X0 X1) = k8_flang_1 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0))) \Rightarrow (k2_flang_2 X0 (k2_flang_2 X0 X1) = k2_flang_2 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k8_afinsq_1 X1))) \Rightarrow ((X0 \in k2_flang_2 X1 X2) \Leftrightarrow ((X0 = k2_flang_1 X1) \vee (X0 \in X2))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k3_catalan2 X0))) \Rightarrow ((k2_flang_1 X0 \in X1) \Rightarrow ((k8_flang_1 X0 X1 = k6_flang_1 X0 (k8_flang_1 X0 X1) X1) \wedge (k8_flang_1 X0 X1 = k6_flang_1 X0 X1 (k8_flang_1 X0 X1)))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0))) \Rightarrow (k6_flang_1 X0 (k2_flang_2 X0 X1) (k8_flang_1 X0 X1) = k6_flang_1 X0 (k8_flang_1 X0 X1) (k2_flang_2 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0. k3_catalan2 X0 = k8_afinsq_1 X0 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0)))\Rightarrow(m1_subset_1 (k2_flang_2 X0 X1) (k1_zfmisc_1 (k8_afinsq_1 X0))) \quad (7)$$

Theorem 1

$$\forall X0.\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0)))\Rightarrow(k6_flang_1 X0 (k2_flang_2 X0 X1) (k8_flang_1 X0 X1) = k8_flang_1 X0 X1)$$