

t109_finseq_6 (TMckjJwsNuauZKDHPm- pAeqH3LNFDDeDBkX)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.\neg(X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((X1 = k9_finseq_1 X0) \Leftrightarrow ((k3_finseq_1 X1 = np_1) \wedge (k10_xtuple_0 X1 = k1_tarski X0))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow \\ & ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.\forall X2.(m2_finseq_1 \\ & X2 X1) \Rightarrow (((r1_xxreal_0 np_1 X0) \wedge (r1_xxreal_0 X0 (k3_finseq_1 \\ & X2))) \Rightarrow (k7_partfun1 X1 X2 X0 = k1_funct_1 X2 X0))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (7)$$

Assume the following.

$$r1_xxreal_0 np_1 np_1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X0) \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (11)$$

Assume the following.

$$\forall X0.k9_finseq_1 X0 = k5_finseq_1 X0 \quad (12)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (13)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (k4_finseq_1 X0 = k9_xtuple_0 X0) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (\\ & k1_relset_1 X0 X1 = k9_xtuple_0 X1) \end{aligned} \quad (15)$$

Assume the following.

$$\forall X0.v1_finseq_1 (k5_finseq_1 X0) \quad (16)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (17)$$

Assume the following.

$$\forall X0.(v1_relat_1 (k5_finseq_1 X0) \wedge (v1_funct_1 (k5_finseq_1 X0))) \quad (18)$$

Assume the following.

$$\forall X0.\exists X1.m1_subset_1 X1 X0 \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge (v1_funct_1 X1) \wedge (v1_finseq_1 X1)) \quad (20)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \wedge ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1)))) \Rightarrow \\ & ((v1_relat_1 (k7_finseq_1 X0 X1)) \wedge ((v1_funct_1 (k7_finseq_1 X0 X1)) \wedge (v1_finseq_1 (k7_finseq_1 X0 X1)))) \quad (21) \end{aligned}$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (22)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \quad (23)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1)))) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge (v1_finseq_1 X2)))) \Rightarrow \\ & ((X2 = k7_finseq_1 X0 X1) \Leftrightarrow ((k4_finseq_1 X2 = k2_finseq_1 (k2_nat_1 (k3_finseq_1 X0) (k3_finseq_1 X1)))) \wedge ((\forall X3.(v7_ordinal1 X3) \Rightarrow ((X3 \in k4_finseq_1 X0) \Rightarrow (k1_funct_1 X2 X3 = k1_funct_1 X0 X3))) \wedge (\forall X3.(v7_ordinal1 X3) \Rightarrow ((X3 \in k4_finseq_1 X1) \Rightarrow (k1_funct_1 X2 (k2_nat_1 (k3_finseq_1 X0) X3) = k1_funct_1 X1 X3)))))) \quad (24) \end{aligned}$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_nat_1 X1 X0) \quad (25)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (26)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow \\ ((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge \\ (v1_finseq_1 X0)))) \quad (27)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (28)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (29)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.\forall X2.(m2_finseq_1 \\ X2 X0)\Rightarrow((r1_xxreal_0 np_1 (k3_finseq_1 X2))\Rightarrow((k1_funct_1 (k7_finseq_1 \\ X2 (k9_finseq_1 X1)) np_1 = k1_funct_1 X2 np_1)\wedge((k1_funct_1 \\ (k7_finseq_1 X2 (k9_finseq_1 X1)) np_1 = k7_partfun1 X0 X2 np_1)\wedge \\ ((k1_funct_1 (k7_finseq_1 (k9_finseq_1 X1) X2) (k2_nat_1 (k3_finseq_1 \\ X2) np_1) = k1_funct_1 X2 (k3_finseq_1 X2))\wedge(k1_funct_1 (k7_finseq_1 \\ (k9_finseq_1 X1) X2) (k2_nat_1 (k3_finseq_1 X2) np_1) = k7_partfun1 \\ X0 X2 (k3_finseq_1 X2))))))))))$$