

t10_autalg_1

(TMH7tmqA7ZnobU1Bk2FFJQYFz6zJGRer3Zo)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_pzfmisc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 X0) \wedge \\ & (v1_funct_1 X1) \wedge (v1_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1_relat_1 \\ & X2) \wedge ((v4_relat_1 X2 X0) \wedge ((v1_funct_1 X2) \wedge (v1_partfun1 X2 X0)))) \Rightarrow \\ & ((r1_pzfmisc1 X0 X1 X2) \Leftrightarrow (\forall X3. ((X3 \in X0) \wedge (k1_funct_1 X2 X3 = \\ & k1_xboole_0))) \Rightarrow (k1_funct_1 X1 X3 = k1_xboole_0))) \end{aligned} \tag{1}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 X0) \wedge \\ & (v1_funct_1 X1) \wedge (v1_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1_relat_1 \\ & X2) \wedge ((v4_relat_1 X2 X0) \wedge ((v1_funct_1 X2) \wedge (v1_partfun1 X2 X0)))) \Rightarrow \\ & (\forall X3. ((v1_relat_1 X3) \wedge ((v4_relat_1 X3 X0) \wedge ((v1_funct_1 \\ & X3) \wedge (v1_partfun1 X3 X0)))) \Rightarrow (((r1_pzfmisc1 X0 X1 X2) \wedge (r1_pzfmisc1 \\ & X0 X2 X3)) \Rightarrow (r1_pzfmisc1 X0 X1 X3))) \end{aligned}$$