

t10_fin_topo (TMR- NUL6miNETAzgVJM2DY3dZ5zV28xB3PrE)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k11_fin_topo : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_fin_topo : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_fin_topo : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 \\ & (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((X1 \in k10_fin_topo X0 X2) \Leftrightarrow ((\\ & X1 \in X2) \wedge (r1_xboole_0 (k7_subset_1 (u1_struct_0 X0) (k1_fin_topo \\ & X0 X1) (k6_domain_1 (u1_struct_0 X0) X1)) X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k7_subset_1 X0 X1 X2 = k4_xboole_0 X1 X2) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k4_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (\neg X3 \in X1))) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (k11_fin_topo \\ & X0 X1 = k7_subset_1 (u1_struct_0 X0) X1 (k10_fin_topo X0 X1))) \end{aligned} \quad (5)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 \\ & (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((X1 \in k11_fin_topo X0 X2) \Leftrightarrow ((\\ & X1 \in X2) \wedge (\neg r1_xboole_0 (k7_subset_1 (u1_struct_0 X0) (k1_fin_topo \\ & X0 X1) (k6_domain_1 (u1_struct_0 X0) X1)) X2)))))) \end{aligned}$$