

t10\_finseq\_6  
(TMZYiopQ3tPt1Rskhyvf4Pbyh7ekNwfWP4W)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1)))) \Rightarrow \\ & ((r1\_tarski X0 X1) \Leftrightarrow (\exists X2.((v1\_relat\_1 X2) \wedge ((v1\_funct\_1 X2) \wedge (v1\_finseq\_1 X2)))) \wedge (X1 = k7\_finseq\_1 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \wedge ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1)))) \Rightarrow \\ & (((v1\_relat\_1 (k7\_finseq\_1 X0 X1)) \wedge ((v1\_funct\_1 (k7\_finseq\_1 X0 X1)) \wedge (v1\_finseq\_1 (k7\_finseq\_1 X0 X1)))) \end{aligned} \quad (2)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1)))) \Rightarrow (r1\_tarski X0 (k7\_finseq\_1 X0 X1)) \end{aligned}$$