

t10_funct_3
(TMU14RZxdiJns2EdzjfX3YfSP1HDybDEPTw)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_3 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (r1_tarski (k10_xtuple_0 (k1_funct_3 X0)) (k1_zfmisc_1 (k10_xtuple_0 X0))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski X1 X2)) \Rightarrow (r1_tarski X0 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (r1_tarski (k7_relat_1 X1 X0) (k10_xtuple_0 X1)) \quad (3)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v1_relat_1 (k1_funct_3 X0)) \wedge (v1_funct_1 (k1_funct_3 X0))) \quad (4)$$

Theorem 1

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (r1_tarski (k7_relat_1 (k1_funct_3 X1) X0) (k1_zfmisc_1 (k10_xtuple_0 X1)))$$