

t10_hilbasis (TMYqZnzHqRHBuRG- zLxh86sLMaePpS2CsXwc)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_hilbasis : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_polynom2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow (k1_polynom2 X0 (k2_hilbasis X0 X1) = k6_domain_1 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k6_domain_1 X0 X1 = k1_tarski X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (3)$$

Theorem 1

$$\forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow (\forall X2. (m1_subset_1 X2 X0) \Rightarrow ((k2_hilbasis X0 X1 = k2_hilbasis X0 X2) \Rightarrow (X1 = X2))))$$