

t10_jordan21 (TMQcCYpCzdvdzMEw- FuTDU5Rq6jGUqUyfm6kr)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_jordan2c : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. r1_xboole_0 (k4_xboole_0 X0 X1) X1 \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_xboole_0 X1 X2)) \Rightarrow (r1_xboole_0 X0 X2) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. (m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 (k15_euclid X0)))) \Rightarrow \\ & (r1_tarski (k2_jordan2c X0 X1) (k3_subset_1 (u1_struct_0 (k15_euclid \\ & X0)) X1))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k3_subset_1 X0 X1 = k4_xboole_0 X0 X1) \quad (5)$$

Theorem 1

$$\forall X0. (m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (r1_xboole_0 (k2_jordan2c np_2 X0) X0)$$