

t10_nat_3

(TMM3jYU3xKzWh3GhVmMuKwB9iYH7nPfrqTB)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k2_nat_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_nat_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k1_newton X0 np_1 = X0) \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 X0 k1_numbers) \wedge (v7_ordinal1 X1)) \Rightarrow (k2_newton X0 X1 = k1_newton X0 X1) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_finseq_1 X0 k1_numbers) \wedge (v7_ordinal1 X1)) \Rightarrow (k2_nat_3 X0 X1 = k1_nat_3 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (7)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(m1_subset_1 (k1_seq_1 X0 X1) k1_numbers) \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v1_finseq_1 X0)\wedge(v3_valued_0 X0))))\Rightarrow(\forall X1.(v7_ordinal1 X1)\Rightarrow(\forall X2. \\ & ((v1_relat_1 X2)\wedge((v1_funct_1 X2)\wedge(v1_finseq_1 X2))))\Rightarrow((X2 = k1_nat_3 X0 X1)\Leftrightarrow((k3_finseq_1 X2 = k3_finseq_1 X0)\wedge(\forall X3. \\ & (X3 \in k4_finseq_1 X2)\Rightarrow(k1_funct_1 X2 X3 = k2_newton (k1_seq_1 X0 X3) X1)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (12)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v5_relat_1 X0 k1_numbers))\Rightarrow((v1_relat_1 X0)\wedge(v3_valued_0 X0)) \quad (13)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xcmplx_0 X0) \quad (14)$$

Assume the following.

$$\forall X0.(v6_membered X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(v7_ordinal1 X1)) \quad (15)$$

Theorem 1 $\forall X0.(m2_finseq_1 X0 k1_numbers)\Rightarrow(k2_nat_3 X0 np_1 = X0).$