

t10_prepower
(TMdQT7HX6bT7425xuuCDMNgc5TrFHCe3w9)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k2_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X1 k6_numbers) \wedge (r1_xxreal_0 (k1_newton X1 X0) k6_numbers))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1)) \quad (3)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((r1_xxreal_0 np_1 X0) \Rightarrow (k2_newton k6_numbers X0 = k6_numbers)) \quad (4)$$

Assume the following.

$$(m2_subset_1 np_0 k1_numbers k5_numbers) \wedge ((m1_subset_1 np_0 k5_numbers) \wedge (m1_subset_1 np_0 k1_numbers)) \quad (5)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (6)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_newton X0 X1 = k1_newton X0 X1) \quad (8)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0)\wedge((v1_xcmplx_0 X0)\wedge((v1_xxreal_0 X0)\wedge(v1_xreal_0 X0))) \quad (9)$$

Assume the following.

$$\begin{aligned} &\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(v1_xreal_0 X1)\Rightarrow(\forall X2. \\ &(v7_ordinal1 X2)\Rightarrow(\neg(\neg r1_xxreal_0 X0 k6_numbers)\wedge((\neg r1_xxreal_0 \\ &X1 X0)\wedge((r1_xxreal_0 np_1 X2)\wedge(r1_xxreal_0 (k1_newton X1 X2) \\ &(k1_newton X0 X2))))))) \end{aligned} \quad (10)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (11)$$

Theorem 1

$$\begin{aligned} &\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(v1_xreal_0 X1)\Rightarrow(\forall X2. \\ &(v7_ordinal1 X2)\Rightarrow(\neg(r1_xxreal_0 k6_numbers X0)\wedge((\neg r1_xxreal_0 \\ &X1 X0)\wedge((r1_xxreal_0 np_1 X2)\wedge(r1_xxreal_0 (k1_newton X1 X2) \\ &(k1_newton X0 X2))))))) \end{aligned}$$