

t10_prgcor_1
(TMNkjRut4Jn9PaTyZ5LQymYxJ3dXtGrZ8YV)

October 27, 2020

Let $v1_int_1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k1_prgcor_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_int_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k3_nat_d : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.(m1_subset_1 \\ X1 k5_numbers) \Rightarrow ((\neg r1_xxreal_0 X1 k6_numbers) \Rightarrow (k1_prgcor_1 X0 \\ X1 = k3_nat_d X0 X1))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.\forall X1. \neg(v1_xboole_0 X0) \wedge ((X0 \neq X1) \wedge (v1_xboole_0 X1)) \tag{2}$$

Assume the following.

$$\forall X0.(v1_int_1 X0) \Rightarrow ((r1_xxreal_0 k6_numbers X0) \Rightarrow (X0 \in k5_numbers)) \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \tag{4}$$

Assume the following.

$$k6_numbers = k1_xboole_0 \tag{5}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{6}$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (k3_nat_d X0 X1 = k5_int_1 X0 X1) \tag{7}$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (8)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (11)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (12)$$

Assume the following.

$$\forall X0.(v1_int_1 X0) \Rightarrow (v1_xreal_0 X0) \quad (13)$$

Theorem 1

$$\forall X0.(v1_int_1 X0) \Rightarrow (\forall X1.(v1_int_1 X1) \Rightarrow ((r1_xxreal_0 k6_numbers X0) \Rightarrow ((r1_xxreal_0 X1 k6_numbers) \vee (k1_prgcor_1 X0 X1 = k5_int_1 X0 X1))))$$