

t10_real.3

(TMHn3RDP25ixz6g2SWv5KaBcAbirv1jJFXF)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $v4_valued_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow (((r1_tarski \\ & (k9_xtuple_0 X2) X0) \wedge (r1_tarski (k10_xtuple_0 X2) X1)) \Rightarrow (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \tag{2}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \tag{4}$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \tag{5}$$

Assume the following.

$$v6_membered k4_ordinal1 \tag{6}$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v4_valued_0 X0)))\Rightarrow(v7_ordinal1 (k1_funct_1 X0 X1)) \quad (7)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow (X2 \in X1)) \quad (9)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.(X1 = k10_xtuple_0 X0)\Leftrightarrow(\forall X2.(X2 \in X1)\Leftrightarrow(\exists X3.(X3 \in k9_xtuple_0 X0)\wedge(X2 = k1_funct_1 X0 X3)))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(((X1 \neq k1_xboole_0)\Rightarrow((v1_funct_2 X2 X0 X1)\Leftrightarrow(X0 = k1_relset_1 X0 X2)))\wedge((X1 = k1_xboole_0)\Rightarrow((v1_funct_2 X2 X0 X1)\Leftrightarrow(X2 = k1_xboole_0)))) \quad (11)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Leftrightarrow(X0 \in k4_ordinal1) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(v6_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v4_valued_0 X2)) \quad (14)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(((v1_funct_1 X0)\wedge((v1_funct_2 X0 k5_numbers k5_numbers)\wedge(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k5_numbers))))))\Leftrightarrow((k9_xtuple_0 X0 = k5_numbers)\wedge(\forall X1.(X1 \in k5_numbers)\Rightarrow(v7_ordinal1 (k1_funct_1 X0 X1))))$$