

# t10\_scmring3

(TMV4jf8WHRunqZg8dKbYxcJYjaptefqXiMw)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_group\_1 : \iota \Rightarrow o$  be given. Let  $v4\_vectsp\_1 : \iota \Rightarrow o$  be given. Let  $v5\_vectsp\_1 : \iota \Rightarrow o$  be given. Let  $l6\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k2\_compos\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $u1\_compos\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_scmring2 : \iota \Rightarrow \iota$  be given. Let  $k8\_scmring2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_6 : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_compos\_0 : \iota \Rightarrow o$  be given. Let  $k4\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $l1\_compos\_1 : \iota \Rightarrow o$  be given. Let  $v2\_compos\_0 : \iota \Rightarrow o$  be given. Let  $v3\_compos\_0 : \iota \Rightarrow o$  be given. Let  $v5\_compos\_0 : \iota \Rightarrow o$  be given. Let  $l1\_extpro\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $l1\_memstr\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_extpro\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_2 : \iota$  be given. Let  $k3\_xtuple\_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k12\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((\neg v1\_xboole\_0 X0) \wedge (v1\_compos\_0 X0)) \wedge \\ & (m1\_subset\_1 X1 X0)) \Rightarrow (k2\_compos\_0 X0 X1 = k4\_xtuple\_0 X1) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k1\_xtuple\_0 (k4\_tarski X0 X1) = X0 \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. (l1\_compos\_1 X0) \Rightarrow ((v1\_compos\_0 (u1\_compos\_1 X0)) \wedge \\ & ((v2\_compos\_0 (u1\_compos\_1 X0)) \wedge ((v3\_compos\_0 (u1\_compos\_1 \\ & X0)) \wedge (v5\_compos\_0 (u1\_compos\_1 X0)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (l1\_extpro\_1 X1 X0) \Rightarrow ((l1\_memstr\_0 X1 X0) \wedge \\ & (l1\_compos\_1 X1)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge \\ & ((v2\_rlvect\_1 X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v3\_group\_1 \\ & X0) \wedge ((v4\_vectsp\_1 X0) \wedge ((v5\_vectsp\_1 X0) \wedge (l6\_algstr\_0 X0)))))))))) \wedge \\ & (m1\_subset\_1 X1 k5\_numbers) \Rightarrow (m1\_subset\_1 (k8\_scmring2 X0 X1) \\ & (u1\_compos\_1 (k1\_scmring2 X0))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\ & X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v3\_group\_1 X0) \wedge \\ & (v4\_vectsp\_1 X0) \wedge ((v5\_vectsp\_1 X0) \wedge (l6\_algstr\_0 X0)))))))))) \Rightarrow \\ & ((v1\_extpro\_1 (k1\_scmring2 X0) np\_2) \wedge (l1\_extpro\_1 (k1\_scmring2 \\ & X0) np\_2)) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\ & X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v3\_group\_1 X0) \wedge \\ & (v4\_vectsp\_1 X0) \wedge ((v5\_vectsp\_1 X0) \wedge (l6\_algstr\_0 X0)))))))))) \Rightarrow \\ & (\forall X1. (m1\_subset\_1 X1 k5\_numbers) \Rightarrow (k8\_scmring2 X0 X1 = k3\_xtuple\_0 \\ & np\_6 (k12\_finseq\_1 k5\_numbers X1) k1\_xboole\_0)) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0. k4\_xtuple\_0 X0 = k1\_xtuple\_0 (k1\_xtuple\_0 X0) \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. k3\_xtuple\_0 X0 X1 X2 = k4\_tarSKI \\ & (k4\_tarSKI X0 X1) X2 \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0. (v5\_compos\_0 X0) \Rightarrow (\neg v1\_xboole\_0 X0) \quad (10)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\ & X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v3\_group\_1 X0) \wedge \\ & (v4\_vectsp\_1 X0) \wedge ((v5\_vectsp\_1 X0) \wedge (l6\_algstr\_0 X0)))))))))) \Rightarrow \\ & (\forall X1. (m1\_subset\_1 X1 k5\_numbers) \Rightarrow (k2\_compos\_0 (u1\_compos\_1 \\ & (k1\_scmring2 X0)) (k8\_scmring2 X0 X1) = np\_6)) \end{aligned}$$