

## t10\_setwiseo

(TMJK1o4LXb2op4Rdm554CHPkuYmDNTA9oyF)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_finsub\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1\_funct\_1 X3) \wedge \\ & ((v1\_funct\_2 X3 X0 X1) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 X1)))) \Rightarrow (\forall X4. \neg (X4 \in k7\_relset\_1 X0 X1 X3 X2) \wedge (\forall X5. \\ & (m1\_subset\_1 X5 X0) \Rightarrow (\neg (X5 \in X2) \wedge (X4 = k1\_funct\_1 X3 X5)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((\neg v1\_xboole\_0 X0) \wedge \\ & (((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))) \wedge (m1\_subset\_1 X3 X0))) \Rightarrow (k3\_funct\_2 X0 \\ & X1 X2 X3 = k1\_funct\_1 X2 X3) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

### Theorem 1

$$\begin{aligned} & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. (\neg v1\_xboole\_0 X1) \Rightarrow \\ & (\forall X2. (m1\_subset\_1 X2 (k5\_finsub\_1 X0)) \Rightarrow (\forall X3. \forall X4. \\ & ((v1\_funct\_1 X4) \wedge ((v1\_funct\_2 X4 X0 X1) \wedge (m1\_subset\_1 X4 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))) \Rightarrow ((\forall X5. (m1\_subset\_1 X5 X0) \Rightarrow ((X5 \in \\ & X2) \Rightarrow (k3\_funct\_2 X0 X1 X4 X5 \in X3)) \Rightarrow (r1\_tarski (k7\_relset\_1 X0 X1 \\ & X4 X2) X3)))))) \end{aligned}$$