

t10_tops_3
 (TMZtzVEytot9sExYie5o3DUmDcg46UvebS4)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v2_tops_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_struct_0 : \iota \Rightarrow \iota$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. ((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc X0))) \Rightarrow (\neg v2_tops_1 (k2_struct_0 X0) X0) \quad (1)$$

Assume the following.

$$\forall X0. (l1_pre_topc X0) \Rightarrow (l1_struct_0 X0) \quad (2)$$

Assume the following.

$$\forall X0. (l1_struct_0 X0) \Rightarrow (k2_struct_0 X0 = u1_struct_0 X0) \quad (3)$$

Theorem 1

$$\forall X0. ((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc X0))) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\neg (v2_tops_1 X1 X0) \wedge (X1 = u1_struct_0 X0)))$$