

t10_waybel14 (TMTA-
jdv2kQKTsiR9RFgEp6iUuSmeSMRVQTh)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v5_orders_2 : \iota \Rightarrow o$ be given. Let $v1_yellow_0 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $k6_waybel_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_yellow_0 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge ((v5_orders_2 X0) \wedge ((v1_yellow_0 X0) \wedge (l1_orders_2 X0)))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (r1_orders_2 X0 (k3_yellow_0 X0) X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow ((\forall X2.(m1_subset_1 X2 X0) \Rightarrow (X2 \in X1)) \Rightarrow (X0 = X1)) \quad (2)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow ((X2 \in k6_waybel_0 X0 X1) \Leftrightarrow (r1_orders_2 X0 X1 X2)))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1.(((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) \wedge (m1_subset_1 X1 (u1_struct_0 X0))) \Rightarrow (m1_subset_1 (k6_waybel_0 X0 X1) (k1_zfmisc_1 (u1_struct_0 X0))) \quad (4)$$

Assume the following.

$$\forall X0.(l1_orders_2 X0) \Rightarrow (m1_subset_1 (k3_yellow_0 X0) (u1_struct_0 X0)) \quad (5)$$

Theorem 1

$$\forall X0.((\neg v2_struct_0 X0) \wedge ((v5_orders_2 X0) \wedge ((v1_yellow_0 X0) \wedge (l1_orders_2 X0)))) \Rightarrow (k6_waybel_0 X0 (k3_yellow_0 X0) = u1_struct_0 X0)$$