

t10\_xxreal\_2  
(TMUA9o7oGVM3NLn72c9HibAcBDDj8oJCukx)

October 27, 2020

Let  $v2\_membered : \iota \Rightarrow o$  be given. Let  $k1\_xxreal\_2 : \iota \Rightarrow \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m1\_xxreal\_2 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.(v2\_membered X0) \Rightarrow (\forall X1.(v2\_membered X1) \Rightarrow (\forall X2. \\ (m1\_xxreal\_2 X2 X0) \Rightarrow (\forall X3.(m1\_xxreal\_2 X3 X1) \Rightarrow (m1\_xxreal\_2 \\ (k4\_xxreal\_0 X2 X3) (k2\_xboole\_0 X0 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.r1\_tarski X0 (k2\_xboole\_0 X0 X1) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(v2\_membered X0) \Rightarrow (\forall X1.(v2\_membered X1) \Rightarrow (( \\ r1\_tarski X0 X1) \Rightarrow (\forall X2.(m1\_xxreal\_2 X2 X1) \Rightarrow (m1\_xxreal\_2 \\ X2 X0)))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ (v1\_xxreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X2 X1)) \Rightarrow \\ (r1\_xxreal\_0 (k4\_xxreal\_0 X0 X2) X1)))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v2\_membered X0) \wedge (v2\_membered X1)) \Rightarrow (v2\_membered (k2\_xboole\_0 X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.(v2\_membered\ X0) \Rightarrow (\forall X1.(m1\_xxreal\_2\ X1\ X0) \Rightarrow (v1\_xxreal\_0\ X1)) \quad (7)$$

Assume the following.

$$\forall X0.(v2\_membered\ X0) \Rightarrow (v1\_xxreal\_0\ (k1\_xxreal\_2\ X0)) \quad (8)$$

Assume the following.

$$\forall X0.(v2\_membered\ X0) \Rightarrow (\forall X1.(v1\_xxreal\_0\ X1) \Rightarrow ((X1 = k1\_xxreal\_2\ X0) \Leftrightarrow ((m1\_xxreal\_2\ X1\ X0) \wedge (\forall X2.(m1\_xxreal\_2\ X2\ X0) \Rightarrow (r1\_xxreal\_0\ X1\ X2)))))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.k2\_xboole\_0\ X0\ X1 = k2\_xboole\_0\ X1\ X0 \quad (10)$$

Assume the following.

$$\forall X0.(v2\_membered\ X0) \Rightarrow (\forall X1.(m1\_subset\_1\ X1\ (k1\_zfmisc\_1\ X0)) \Rightarrow (v2\_membered\ X1)) \quad (11)$$

**Theorem 1**

$$\forall X0.(v2\_membered\ X0) \Rightarrow (\forall X1.(v2\_membered\ X1) \Rightarrow (k1\_xxreal\_2\ (k2\_xboole\_0\ X0\ X1) = k4\_xxreal\_0\ (k1\_xxreal\_2\ X0)\ (k1\_xxreal\_2\ X1)))$$