

# t111\_flgang\_2 (TMZkZvTmF- sJLJguA5bMied3TxQsbMDqCjMB)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k8\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k6\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_flang\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_flang\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow (k2\_flang\_2 X0 X1 = k1\_flang\_2 X0 X1 k6\_numbers np\_1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow (\forall X2. (v7\_ordinal1 X2) \Rightarrow (\forall X3. (v7\_ordinal1 X3) \Rightarrow (\forall X4. (v7\_ordinal1 X4) \Rightarrow (\forall X5. (v7\_ordinal1 X5) \Rightarrow \\ & (k6\_flang\_1 X0 (k1\_flang\_2 X0 X1 X2 X3) (k1\_flang\_2 X0 X1 X4 X5) = k6\_flang\_1 X0 (k1\_flang\_2 X0 X1 X4 X5) (k1\_flang\_2 X0 X1 X2 X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$m1\_subset\_1 k1\_xboole\_0 k4\_ordinal1 \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (5)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0. (m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (7)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 \\ & \quad X0))) \Rightarrow (\forall X2. (v7\_ordinal1 X2) \Rightarrow (\forall X3. (v7\_ordinal1 \\ X3) \Rightarrow (k6\_flang\_1 X0 (k2\_flang\_2 X0 X1) (k1\_flang\_2 X0 X1 X2 X3) = k6\_flang\_1 \\ & \quad X0 (k1\_flang\_2 X0 X1 X2 X3) (k2\_flang\_2 X0 X1)))) \end{aligned}$$