

t111_xcplx_1
(TMctbSxSH38cDDiS3C7pkY5YszJM1b34Gw7)

October 27, 2020

Let $v1_xcplx_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k7_xcplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_xcplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (k7_xcplx_0 k6_numbers X0 = k6_numbers) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow (\neg (X0 \neq k6_numbers) \wedge ((X1 \neq k6_numbers) \wedge (k7_xcplx_0 X0 X1 = k6_numbers)))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (X1 = k7_xcplx_0 (k3_xcplx_0 X1 X0) X0))) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (k7_xcplx_0 X0 X0 = np_1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcplx_0 X0) \wedge (v1_xcplx_0 X1)) \Rightarrow (v1_xcplx_0 (k7_xcplx_0 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcplx_0 X0) \wedge (v1_xcplx_0 X1)) \Rightarrow (v1_xcplx_0 (k3_xcplx_0 X0 X1)) \quad (6)$$

Theorem 1

$$\forall X0.(v1_xcplx_0 X0) \Rightarrow (\forall X1.(v1_xcplx_0 X1) \Rightarrow (\neg (X0 \neq k6_numbers) \wedge ((X1 \neq k6_numbers) \wedge (k7_xcplx_0 np_1 (k3_xcplx_0 X0 X1) = k6_numbers))))$$