

t112_jgraph_4

(TMZXVrRVwiGiUv9Qs5Qgrv2piL9H5NXUUe5)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k4_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k7_jgraph_4 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k13_complex1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $k12_euclid : \iota \Rightarrow \iota$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_square_1 : \iota \Rightarrow \iota$ be given. Let $k9_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k4_square_1 : \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k16_euclid : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\
 & (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X1 (k13_complex1 \\
 & (k18_euclid X0) (k12_euclid X0))) \Rightarrow ((r1_xxreal_0 (k17_euclid \\
 & X0) k6_numbers) \vee (k3_funct_2 (u1_struct_0 (k15_euclid np_2)) \\
 & (u1_struct_0 (k15_euclid np_2)) (k7_jgraph_4 X1) X0 = k19_euclid \\
 & (k8_real_1 (k12_euclid X0) (k7_square_1 (k9_real_1 np_1 (k4_square_1 \\
 & (k13_complex1 (k6_xcmplx_0 (k13_complex1 (k18_euclid X0) (k12_euclid \\
 & X0)) X1) (k9_real_1 np_1 X1)))))) (k8_real_1 (k12_euclid X0) (\\
 & k13_complex1 (k6_xcmplx_0 (k13_complex1 (k18_euclid X0) (k12_euclid \\
 & X0)) X1) (k9_real_1 np_1 X1)))))) \wedge ((r1_xxreal_0 (k17_euclid \\
 & X0) k6_numbers) \Rightarrow (k3_funct_2 (u1_struct_0 (k15_euclid np_2)) \\
 & (u1_struct_0 (k15_euclid np_2)) (k7_jgraph_4 X1) X0 = X0))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \tag{2}$$

Assume the following.

$$k4_struct_0 (k15_euclid np_2) = k19_euclid k6_numbers k6_numbers \tag{3}$$

Assume the following.

$$(k17_euclid (k4_struct_0 (k15_euclid np_2)) = k6_numbers) \wedge (k18_euclid (k4_struct_0 (k15_euclid np_2)) = k6_numbers) \quad (4)$$

Assume the following.

$$(m2_subset_1 np_0 k1_numbers k5_numbers) \wedge ((m1_subset_1 np_0 k5_numbers) \wedge (m1_subset_1 np_0 k1_numbers)) \quad (5)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (6)$$

Assume the following.

$$r1_xxreal_0 np_0 np_0 \quad (7)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (8)$$

Assume the following.

$$k16_euclid np_2 = k4_struct_0 (k15_euclid np_2) \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k19_euclid X0 X1) (u1_struct_0 (k15_euclid np_2))) \quad (10)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (11)$$

Theorem 1

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (k4_struct_0 (k15_euclid np_2) = k3_funct_2 (u1_struct_0 (k15_euclid np_2)) (u1_struct_0 (k15_euclid np_2)) (k7_jgraph_4 X0) (k4_struct_0 (k15_euclid np_2)))$$