

t114_flang_2 (TMXNWn-
QkaYb7mMhwAFcKNTRwPxXKo3Ahy1R)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_flang_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_flang_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & \quad X0))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (\forall X3. (v7_ordinal1 \\ & \quad X3) \Rightarrow (r1_tarski (k1_flang_2 X0 X1 X2 X3) (k8_flang_1 X0 X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & \quad X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k8_afinsq_1 \\ & \quad X0))) \Rightarrow ((r1_tarski X1 (k8_flang_1 X0 X2)) \Rightarrow (r1_tarski (k2_flang_2 \\ & \quad X0 X1) (k8_flang_1 X0 X2)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((m1_subset_1 X1 \\ & \quad (k1_zfmisc_1 (k8_afinsq_1 X0))) \wedge ((v7_ordinal1 X2) \wedge (v7_ordinal1 \\ & \quad X3))) \Rightarrow (m1_subset_1 (k1_flang_2 X0 X1 X2 X3) (k1_zfmisc_1 (k8_afinsq_1 \\ & \quad X0))) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & \quad X0))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (\forall X3. (v7_ordinal1 \\ & \quad X3) \Rightarrow (r1_tarski (k2_flang_2 X0 (k1_flang_2 X0 X1 X2 X3)) (k8_flang_1 \\ & \quad X0 X1)))) \end{aligned}$$