

t114_funct_7

(TMUei6bUfMonWa48LMMd2z1TS2knJnEUzCt)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & k1_funct_1 (k1_funct_4 X0 (k16_funcop_1 X1 X2)) X1 = X2) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & \forall X3. \forall X4. \forall X5. \neg (X5 \neq X4) \wedge ((X5 \neq X1) \wedge (k1_funct_1 \\ & (k1_funct_4 (k1_funct_4 X0 (k16_funcop_1 X1 X2)) (k16_funcop_1 \\ & X4 X3)) X5 \neq k1_funct_1 X0 X5))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (v1_relat_1 (k16_funcop_1 X0 X1)) \wedge (v1_funct_1 \\ & (k16_funcop_1 X0 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \wedge ((\\ & v1_relat_1 X1) \wedge (v1_funct_1 X1))) \Rightarrow ((v1_relat_1 (k1_funct_4 X0 \\ & X1)) \wedge (v1_funct_1 (k1_funct_4 X0 X1))) \end{aligned} \quad (4)$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & \forall X3. \forall X4. \forall X5. \forall X6. \neg (X1 \neq X3) \wedge ((X1 \neq X5) \wedge \\ & (k1_funct_1 (k1_funct_4 (k1_funct_4 (k1_funct_4 X0 (k16_funcop_1 \\ & X1 X2)) (k16_funcop_1 X3 X4)) (k16_funcop_1 X5 X6)) X1 \neq X2))) \end{aligned}$$