

t114_rvsum_1 (TM-
cjCXfMXqpK3zcH89G8KWgY8xLyEVTSaCm)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k6_rvsum_1 : \iota \Rightarrow \iota$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k30_valued_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ & ((k9_xtuple_0 (k30_valued_1 X0) = k9_xtuple_0 X0) \wedge (\forall X1. \\ & k1_funct_1 (k30_valued_1 X0) X1 = k4_xcmplx_0 (k1_funct_1 X0 X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1))) \Rightarrow ((k3_finseq_1 X0 = k3_finseq_1 X1) \Leftrightarrow (k1_relset_1 k5_numbers \\ & X0 = k1_relset_1 k5_numbers X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (k6_rvsum_1 X0 = k30_valued_1 X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1.((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v1_valued_0 \\ & X0) \wedge (v1_finseq_1 X0)))) \Rightarrow ((v1_relat_1 (k30_valued_1 X0)) \wedge ((\\ & v1_funct_1 (k30_valued_1 X0)) \wedge ((v1_valued_0 (k30_valued_1 X0)) \wedge \\ & (v1_finseq_1 (k30_valued_1 X0)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \Rightarrow ((v1_relat_1 X0) \wedge (v1_valued_0 X0)) \quad (6)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 \ k5_numbers) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0)))) \quad (7)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (k3_finseq_1 (k6_rvsum_1 X0) = k3_finseq_1 X0)$$