

t115\_funct\_4 (TM-  
Rvw7LbxJuazSJcqkQCNfAJxUNs8oaAwVw)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $r1\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_xboole\_0 (k9\_xtuple\_0 \\ X0) (k9\_xtuple\_0 X1)) \Rightarrow (k1\_funct\_4 X0 X1 = k1\_funct\_4 X1 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (\forall X2.((v1\_relat\_1 X2) \wedge \\ (v1\_funct\_1 X2)) \Rightarrow (k1\_funct\_4 (k1\_funct\_4 X0 X1) X2 = k1\_funct\_4 \\ X0 (k1\_funct\_4 X1 X2)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1.(((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \wedge (( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1))) \Rightarrow (k1\_funct\_4 X0 X0 = X0) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1.(((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \wedge (( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1))) \Rightarrow ((v1\_relat\_1 (k1\_funct\_4 X0 \\ X1)) \wedge (v1\_funct\_1 (k1\_funct\_4 X0 X1))) \end{aligned} \quad (4)$$

**Theorem 1**

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (\forall X2.((v1\_relat\_1 X2) \wedge \\ (v1\_funct\_1 X2)) \Rightarrow ((r1\_xboole\_0 (k9\_xtuple\_0 X0) (k9\_xtuple\_0 \\ X1)) \Rightarrow (k1\_funct\_4 (k1\_funct\_4 (k1\_funct\_4 X2 X0) X1) X0 = k1\_funct\_4 \\ (k1\_funct\_4 X2 X0) X1)))) \end{aligned}$$