

t115_zf_lang1

(TMG316Wosy7RwPrW5Z3eeFNF4uLYBsE1ciJ)

October 27, 2020

Let $v1_zf_lang : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zf_lang : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_zf_model : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k11_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_zf_model : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & ((v1_zf_lang X1) \wedge (m2_finseq_1 X1 k5_numbers)) \Rightarrow (\forall X2. (\\ & \neg v1_xboole_0 X2) \Rightarrow (\forall X3. ((v1_funct_1 X3) \wedge ((v1_funct_2 \\ & X3 k1_zf_lang X2) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 k1_zf_lang \\ & X2)))))) \Rightarrow ((r1_zf_model X2 X3 (k11_zf_lang X0 (k10_zf_lang X0 X1)))) \wedge \\ & (r2_zf_model X2 (k11_zf_lang X0 (k10_zf_lang X0 X1)))))) \end{aligned} \quad (1)$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & (\neg v1_xboole_0 X1) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 k1_zf_lang X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k1_zf_lang \\ & X1)))))) \Rightarrow ((r1_zf_model X1 X2 (k11_zf_lang X0 (k10_zf_lang X0 X0))) \wedge \\ & (r2_zf_model X1 (k11_zf_lang X0 (k10_zf_lang X0 X0)))))) \end{aligned}$$