

t118_xreal_1
(TMTinRxJ3DeN64VZrDZhtJTV8Cj5oqAfUtg)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $c7_xreal_0 : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (v2_xxreal_0 X0)) \Rightarrow (v2_xxreal_0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow ((r1_xxreal_0 (k3_xcmplx_0 X2 X1) (k3_xcmplx_0 X3 X0)) \Rightarrow ((r1_xxreal_0 X0 k6_numbers) \vee ((r1_xxreal_0 X1 k6_numbers) \vee (r1_xxreal_0 (k7_xcmplx_0 X2 X0) (k7_xcmplx_0 X3 X1)))))))))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 k6_numbers X2)) \Rightarrow (r1_xxreal_0 (k3_xcmplx_0 X0 X2) (k3_xcmplx_0 X1 X2)))))) \quad (3)$$

Assume the following.

$$c7_xreal_0 = k6_numbers \quad (4)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow ((v2_xxreal_0 X0) \Leftrightarrow (\neg r1_xxreal_0 X0 k6_numbers)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k3_xcmplx_0 X0 X1 = k3_xcmplx_0 X1 X0) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 k6_numbers X1) \wedge (r1_xxreal_0 \\ & X0 X2)) \Rightarrow ((r1_xxreal_0 X0 k6_numbers) \vee (r1_xxreal_0 (k7_xcmplx_0 \\ & X1 X2) (k7_xcmplx_0 X1 X0)))))) \end{aligned}$$