

t119_funct_4 (TMRqYKWxSNXPDhrkxyxoMZf- bLdebtQm3U51)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (r1_xboole_0 X0 X1) \Rightarrow (k4_xboole_0 (k2_xboole_0 X0 X1) X1 = X0) \quad (1)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_xboole_0 (k9_xtuple_0 X0) (k9_xtuple_0 X1)) \Rightarrow (k2_xboole_0 X0 X1 = k1_funct_4 X0 X1))) \quad (2)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (\forall X1. (v1_relat_1 X1) \Rightarrow ((r1_xboole_0 (k9_xtuple_0 X0) (k9_xtuple_0 X1)) \Rightarrow (r1_xboole_0 X0 X1))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (4)$$

Theorem 1

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_xboole_0 (k9_xtuple_0 X0) (k9_xtuple_0 X1)) \Rightarrow (k6_subset_1 (k1_funct_4 X0 X1) X1 = X0)))$$