

t11_cfunct_1

(TMVXg3H2ujp4CAQeWTF49Cb9BKfQuLfgkgr)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_numbers : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_cfunct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarSKI : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarSKI : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_complex1 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \Rightarrow (\\ k8_relset_1 X0 k2_numbers (k2_cfunct_1 X0 X1) (k1_tarSKI k6_numbers) = \\ k1_xboole_0)) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k9_xtuple_0 (k5_relat_1 X1 X0) = k3_xboole_0 (k9_xtuple_0 X1) X0) \tag{2}$$

Assume the following.

$$\forall X0. k4_xboole_0 X0 k1_xboole_0 = X0 \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. r1_tarSKI (k4_xboole_0 X0 X1) X0 \tag{4}$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarSKI X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(k8_relset_1 X0 X1 X2 X3 = k8_relat_1 X2 X3) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(k7_subset_1 X0 X1 X2 = k4_xboole_0 X1 X2) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow(k2_partfun1 X0 X1 X2 X3 = k5_relat_1 X2 X3) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X0)\Rightarrow(v1_relat_1 (k5_relat_1 X0 X1)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow(((v1_funct_1 (k2_partfun1 X0 X1 X2 X3))\wedge(m1_subset_1 (k2_partfun1 X0 X1 X2 X3) (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers))))))\Rightarrow(((v1_funct_1 (k2_cfunct_1 X0 X1))\wedge(m1_subset_1 (k2_cfunct_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(m1_subset_1 (k1_relset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (13)$$

Assume the following.

$$k1_xboole_0 = the (\lambda X0 : \iota.v1_xboole_0 X0) \quad (14)$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \Rightarrow (\\ \forall X2.((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ X0 k2_numbers)))) \Rightarrow ((X2 = k2_cfunct_1 X0 X1) \Leftrightarrow ((k1_relset_1 X0 X2 = \\ k7_subset_1 X0 (k1_relset_1 X0 X1) (k8_relset_1 X0 k2_numbers X1 \\ (k1_tarSKI k6_numbers))) \wedge (\forall X3.(m1_subset_1 X3 X0) \Rightarrow ((\\ X3 \in k1_relset_1 X0 X2) \Rightarrow (k7_partfun1 k2_numbers X2 X3 = k12_complex1 \\ (k7_partfun1 k2_numbers X1 X3)))))))))) \end{aligned} \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (18)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \Rightarrow (\\ k1_relset_1 X0 (k2_cfunct_1 X0 (k2_cfunct_1 X0 X1)) = k1_relset_1 \\ X0 (k2_partfun1 X0 k2_numbers X1 (k1_relset_1 X0 (k2_cfunct_1 X0 \\ X1)))))) \end{aligned}$$