

t11_cqc_lang
(TMJbz3DAd948wmY8aBeSkdYG4z88JqjQ1zb)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $k3_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k14_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k13_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0) \Rightarrow (\forall X2.(m1_subset_1 X2 (k9_qc_lang1 X0) \Rightarrow ((m2_subset_1 \\ (k14_qc_lang1 X0 X1 X2) (k9_qc_lang1 X0) (k3_cqc_lang X0)) \Leftrightarrow ((m2_subset_1 \\ X1 (k9_qc_lang1 X0) (k3_cqc_lang X0)) \wedge (m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0))))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0) \Rightarrow ((m2_subset_1 (k13_qc_lang1 X0 X1) (k9_qc_lang1 X0) (k3_cqc_lang \\ X0)) \Leftrightarrow (m2_subset_1 X1 (k9_qc_lang1 X0) (k3_cqc_lang X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\neg v1_xboole_0 (k3_cqc_lang X0)) \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2 X0 X1) \Rightarrow (m1_subset_1 X2 X0)) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (m1_subset_1 (k3_cqc_lang X0) (k1_zfmisc_1 \\ (k9_qc_lang1 X0))) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k9_qc_lang1 \\ & X0)) \Rightarrow (\forall X2.(m1_subset_1\ X2\ (k9_qc_lang1\ X0)) \Rightarrow (k3_qc_lang2 \\ & X0\ X1\ X2 = k13_qc_lang1\ X0\ (k14_qc_lang1\ X0\ (k13_qc_lang1\ X0\ X1)\ (\\ & k13_qc_lang1\ X0\ X2)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ X0)) \Rightarrow (v1_xboole_0\ X1)) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ & X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ & X0)\ (k3_cqc_lang\ X0)) \Rightarrow (m2_subset_1\ (k3_qc_lang2\ X0\ X1\ X2)\ (k9_qc_lang1 \\ & X0)\ (k3_cqc_lang\ X0)))) \end{aligned}$$