

t11_euclid_3 (TMRCsHJfafLSoYmRip- nEuA7KKTHLc7Y9Fc3)

October 27, 2020

Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k1_euclid_3 : \iota \Rightarrow \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k4_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k3_complex1 : \iota \Rightarrow \iota$ be given. Let $k4_complex1 : \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ (k4_algstr_0 (k15_euclid np_2) X0 = k19_euclid (k1_real_1 (k17_euclid \\ X0)) (k1_real_1 (k18_euclid X0))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid \\ (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xcmplx_0 X0) \Rightarrow (k19_euclid (k3_complex1 (k4_xcmplx_0 \\ X0)) (k4_complex1 (k4_xcmplx_0 X0)) = k19_euclid (k1_real_1 (k3_complex1 \\ X0)) (k1_real_1 (k4_complex1 X0))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (v1_xcmplx_0 (k4_xcmplx_0 X0)) \tag{4}$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (m1_subset_1 (k4_complex1 X0) k1_numbers) \tag{5}$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (m1_subset_1 (k3_complex1 X0) k1_numbers) \tag{6}$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (m1_subset_1 (k1_euclid_3 X0) (u1_struct_0 (k15_euclid np_2))) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k1_euclid_3 X0 = k19_euclid (k3_complex1 X0) (k4_complex1 X0)) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (9)$$

Theorem 1

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k1_euclid_3 (k4_xcmplx_0 X0) = k4_algstr_0 (k15_euclid np_2) (k1_euclid_3 X0))$$