

t11_flang_2
(TMGZVYV6zfGeiLVeT8R6dtZQ6XJbY9rV7b8)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k7_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_catalan2 : \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k3_catalan2 \\ & \quad X0))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (\forall X3. (v7_ordinal1 \\ & X3) \Rightarrow (k7_flang_1 X0 (k7_flang_1 X0 X1 X2) X3 = k7_flang_1 X0 X1 (k3_xcmplx_0 \\ & \quad X2 X3)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. k3_catalan2 X0 = k8_afinsq_1 X0 \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k3_xcmplx_0 X0 X1 = k3_xcmplx_0 X1 X0) \tag{3}$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \tag{4}$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \tag{5}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 \\ & \quad X0))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (\forall X3. (v7_ordinal1 \\ & X3) \Rightarrow (k7_flang_1 X0 (k7_flang_1 X0 X1 X2) X3 = k7_flang_1 X0 (k7_flang_1 \\ & \quad X0 X1 X3) X2))) \end{aligned}$$