

# t11\_grfunc\_1 (TMNzpRb- HEpeu7Hp24KMLSpaRFAHwnWjULPQ)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k3\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarski X0 X1) \Leftrightarrow ((r1\_tarski \\ (k9\_xtuple\_0 X0) (k9\_xtuple\_0 X1)) \wedge (\forall X2.(X2 \in k9\_xtuple\_0 \\ X0) \Rightarrow (k1\_funct\_1 X0 X2 = k1\_funct\_1 X1 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(r1\_tarski \\ X1 X0) \Rightarrow ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. r1\_tarski (k3\_xboole\_0 X0 X1) X0 \quad (3)$$

## Theorem 1

$$\begin{aligned} \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (\forall X2. \\ ((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow ((X0 \in k9\_xtuple\_0 (k3\_xboole\_0 \\ X1 X2)) \Rightarrow (k1\_funct\_1 (k3\_xboole\_0 X1 X2) X0 = k1\_funct\_1 X1 X0))) \end{aligned}$$